


12.25 in

NALOXONE

OVERDOSE EMERGENCY KIT


OPEN LATCH →



HOW TO RECOGNIZE AN OVERDOSE

The following are signs and symptoms to look for to identify a potential overdose:

- Loss of consciousness, limp body
- Slowed breathing, choking
- Small, "pinpoint" pupils
- Pale, blue or cold skin
- Slowed heart rate



Instructions
Inside Cabinet

In An Emergency
Dial 911


CLIENT INFO

Free Naloxone - Be Prepared


OPEN LATCH →

WARNING:
Naloxone freezes at temperatures below 5°F (-15°C). If this happens, the device will not spray. If naloxone is frozen and is needed in an emergency, do NOT wait for the naloxone to thaw. Get emergency medical help right away.

BEWARE OF COUNTERFEIT DRUGS THAT MAY CONTAIN FENTANYL



16.25 in



How to use Naloxone Nasal Spray.


Before use, check for signs of opioid overdose:

- Will not wake up or respond to your voice or touch
- Breathing is very slow, irregular, or has stopped
- Center part of their eye is very small, sometimes called "pinpoint pupils"

Lay the person on their back to receive a dose of Naloxone Nasal Spray.


- 1**

1. **Peel** back the package to remove the device. Hold the device with your thumb on the bottom of the plunger and 2 fingers on the nozzle.


- 2**


2. **Place** and hold the tip of the nozzle in either nostril until your fingers touch the bottom of the patient's nose.

Tilt the person's head back and provide support under the neck with your hand. Gently insert the tip of the nozzle into one nostril, until your fingers on either side of the nozzle are against the bottom of the person's nose.


- 3**


3. **Press** the plunger firmly to give the dose of Naloxone Nasal Spray into the patient's nose.

Remove the Naloxone Nasal Spray from the nostril after giving the dose.



Move the person on their side (recovery position) after giving Naloxone Nasal Spray.

HAND SUPPORTS HEAD



KNEE STOPS BODY FROM ROLLING ONTO STOMACH
- 4**

4. **Get emergency medical help right away.**


 - Watch the person closely. If the person does not respond by waking up, to voice or touch, or breathing normally another dose may be given.
 - Naloxone Nasal Spray may be dosed every 2 to 3 minutes, if available.
 - Repeat Step 2 using a new Naloxone Nasal Spray to give another dose in the other nostril.
 - If additional Naloxone Nasal Sprays are available, repeat step 2 every 2 to 3 minutes until the person responds or emergency medical help is received.

Use Naloxone Nasal Spray (naloxone hydrochloride) for known or suspected opioid overdose in adults and children.
Important: For use in the nose only. Do not remove or test the Naloxone Nasal Spray until ready to use.

Illinois Supply Company
CREATORS OF AEK BEARD STORAGE SYSTEMS
EPIPHORNE - NALOXONE - BEHAVIOR CONTROL

IN AN EMERGENCY TAKE INSTRUCTIONS WITH YOU

PULL
↓



11.75 in

CLIENT INFO

custom vinyl sticker made inhouse (fast)

3 in

